When wheezing is a problem..

.CAM clears the airways.

 synergistic combination to provide maximum bronchodilation with minimal side-effects

complementary action: ephedrine:

relaxes smooth bronchial muscle stimulates respiratory centre butethamate:

opposes bronchoconstriction reduces secretion

CAM bronchial antispasmodic mixture.

- eases sputum raising
- bedtime dose controls nocturnal coughing and wheezing
- suitable for infants, children and adults (except those with severe heart disease or hypertension)

Rybar

RYBAR LABORATORIES LTD., Amersham, England.



PRESCRIBING INFORMATION Presentation: White, sugar-coated tablets each containing 135 mg mebeverine hydrochloride. Available in packs of 100. Basic NHS price £8.35. Yellow, banana-flavoured sugar-free suspension containing mebeverine pamoate equivalent to 50 mg mebeverine hydrochloride per 5 ml. Available in bottles of 300 ml. Basic NHS price £3.50. Indications: 1. Irritable bowel syndrome. 2. Gastrointestinal spasm secondary to organic diseases. Dosage and Administration: Tablets: Adults and children ten years and over: One tablet three times a day, preferably 20 minutes before meals. Suspension: Adults and children ten years and over: 15 ml (150 mg) three times a day, preferably 20 minutes before meals. Contra-indications, warnings, etc: Animal experiments have failed to show any teratogenic effects. However, the usual precautions concerning the administration of any drug during pregnancy should be observed. **Product Licence Number:** Tablets: 0512/0044; Suspension: 0512/0061. Further information is available on request to the Company.

duphar Duphar Laboratories Limited, Gaters Hill, West End, Southampton, SO3 3JD. Telephone: 0703 472281

Write one

Write effect

Write long tern



PRESCRIBING NOTES FOR 'TENORMIN'

HYPERTENSION + ANGINA PROPHYLAXIS

Tenormin' - 100mg atenolol orally once a day.

CHILDREN - There is no experience with 'Tenormin' in children.

ELDERLY PATIENTS -CONTRA-INDICATIONS - Dosage requirements may be lower in elderly patients.

Heart Block, cardiogenic shock.

PRECAUTIONS

PROCAUTIONS
Untreated cardiac failure, bradycardia, renal failure, anaesthesia, pregnancy. Caution in patients with chronic obstructive airways disease or asthma. Atenolol modifies the tachycardia of hypoglycaemia. Co-administration with verapamii or Class I antidysrhythmic agents. Withdrawal of clonidine. Withdrawal of beta blocking drugs should be gradual in patients with ischaemic heart disease.

SIDE-EFFECTS


Coldness of extremities, bradycardia and muscular fatigue may occur. Sleep disturbance rarely seen. Rashes and dry eyes have been reported with beta blockers – consider

discontinuance if they occur.

PRODUCT LICENCE NUMBER AND BASIC NHS COST:
Tenormin Tablets 29 0122 in calendar packs of 28, £6,98.

Tenormin' is a trade mark.

Further information is available on request from the Company



'HBISCRUB'

Effective hand hygiene that persists

'HBISOL'

Hand hygiene on the move

'HIBISCRUB' and 'HIBISOL' Your partners in infection control

'HIBISCRUB' and 'HIBISOL' are trade marks




ICI ANTISEPTICS - BREAKING THE CHAIN OF INFECTION



Further information available on request:

ICI Pharmaceuticals (UK), Alderley House, Alderley Park, Macclesfield, Cheshire, SK10 4TF



BEECHAM RESEARCH

Further information on Augmentin, a product of British research, is available on request to Beecham Research Laboratories, Brentford, Middx TW8 9BD.

Augmentin is a trademark